*TABLE 1: Characteristics of Michigan Residents Living with HIV or AIDS

.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	TABLE 1: Characteristics of Michigan Residents Living with filt of AiDS								
	Estimate of HIV	Estimated	Reported	_	Reported Living with				
	Prevalence ¹	Prevalence	with AIDS ³		HIV not AIDS ³				
	Trevalence	Rate ²	Number	Percent ⁴		Percent⁴			
MICHIGAN TOTAL	15,500	156	5,135	100%	5,581	100%			
SEX	·								
Male	11,940	245	4,109	80%	4,166	75%			
Female	3,570	70	1,026	20%	1,415	25%			
BEHAVIOR	,		,		,				
Male-Male Sex	8,530	N/A	2512	57%	2332	53%			
Injecting Drug Use ⁵	2,950	N/A	881	20%	796	18%			
IDU w/ heterosexual	1,400	N/A	416	9%	379	9%			
IDU w/o heterosexual	1,550	N/A	465	10%	417	9%			
Male-Male Sex/IDU	930	N/A	282	6%	279	6%			
Blood Products	160	N/A	76	2%	50	1%			
Heterosexual ⁶	2,640	N/A	645	15%		19%			
Partner IDU	780	N/A	207	5%	265	6%			
Partner Bisexual	160	N/A	28	1%	40	1%			
Partner Rec'd Bld	60	N/A	13	0%	22	1%			
Partner HIV +	1,550	N/A	397	9%	516	12%			
Perinatal	160	N/A	33	1%	98	2%			
Undetermined/Other ⁴	Not Applicable	N/A	706	(14%)	1183	(21%)			
Presumed Hetersexual ⁷	Not Applicable	N/A	548	(11%)	789	(14%)			
Other ⁸	Not Applicable	N/A	158	(3%)	394	(7%)			
AGE AT DIAGNOSIS						· , ,			
0 -12 years	160	9	34	1%	114	2%			
13 -19 years	310	31	44	1%	184	3%			
20 -24 years	1,400	217	232	5%	725	13%			
25 -29 years	2,480	379	614	12%		19%			
30 -34 years	3,260	461	1095	21%	1132	20%			
35 -39 years	3,100	394	1149	22%	1013	18%			
40 -44 years	2,330	287	918	18%	647	12%			
45 -49 years	1,240	169	552	11%	350	6%			
50 -54 years	780	123	303	6%	201	4%			
55 -59 years	310	64	114	2%	87	2%			
60 -64 years	160	42	52	1%	37	1%			
65 years and over	70	6	28	1%	19	0%			
RACE / ETHNICITY				<u> </u>					
White, Non-Hisp.	5,740	74	1,984	39%		36%			
Black, Non-Hisp.	7,600	542	2,911	57%		60%			
Hispanic	620	191	206	4%		3%			
Asian	50	28	23	0%	14	0%			
American Indian	50	94	10	0%	26	0%			
Unspecified/Other ⁴	Not Applicable	N/A	1	(0%)	136	(2%)			

^{*} See Technical Notes for footnotes to Table 1.

TABLE 2: Michigan Residents Reported Living with HIV or AIDS by County, as of 4/1/03 and Residents Ever Diagnosed with AIDS, 1981 to Present

and Residents Ever Diagnosed with AIDS, 1981 to Present								
Residence at Time Estimate of HIV		Estimated Persons		Persons	Persons Ever			
of Diagnosis	Prevalence ¹	Prevalence		Living with	Diagnosed			
		Rate ²	AIDS ³	HIV not AIDS ³	with AIDS			
Total Michigan	15,500	156.0	5,135	5,581	12,760			
Alcona	10		1	0	3			
Alger	10		1	0	1			
Allegan	90	85.2	39	22	78			
Alpena	10		1	2	8			
Antrim	10		2	1	6			
Arenac	10		0	2	2			
Baraga	10		4	3	8			
Barry	30	52.9	11	6	23			
Bay	80	72.6	20	30	52			
Benzie	10		0	1	2			
Berrien	260	160.0	83	82	170			
Branch	10		1	8	13			
Calhoun	150	108.7	38	58	96			
Cass	30	58.7	8	13	17			
Charlevoix	10		3	6	5			
Cheboygan	10		2	0	5			
Chippewa	10	25.9	2	6	4			
Clare	10	32.0	3	6	8			
Clinton	40	61.8	17	9	28			
Crawford	10		2	0	4			
Delta	20	51.9	6	7	7			
Dickinson	10		2	2	5			
Eaton	40	38.6	14	14	32			
Emmet	10		5	3	9			
Genesee	600	137.6	171	217	410			
Gladwin	10		3	1	5			
Gogebic	10		1	3	7			
Grand Traverse	60	77.3	17	23	35			
Gratiot	10		2	3	8			
Hillsdale	10		2	5	14			
Houghton	10		4	3	8			
Huron	10		2	0	8			
Ingham	430	153.8	119	160	329			
Ionia	20	32.5	9	4	18			
losco	10		1	2	2			
Iron	10		0	1	2			
Isabella	20	31.6	7	6	18			
Jackson	140	88.3	41	50	94			
Kalamazoo	330	138.3	110	101	232			
Kalkaska	10		1	3	4			
Kent	790	137.6	250	262	603			
Keweenaw	10		0	0	0			
Lake	10		6	3	10			
Lapeer	20	22.8	4	11	13			
Leelanau	10		5	0	13			
Lenawee	60	60.7	19	22	41			
Livingston	40	25.5	12	14	31			
Luce	10		0	0	1			

TABLE 2 continued: Michigan Residents Living with HIV or AIDS by County, as of 4/1/03 and Residents Ever Diagnosed with AIDS, 1981 to Present

and Residents Ever Diagnosed with AIDS, 1981 to Present									
Residence at Time	Estimate of HIV	Estimated	Persons	Persons	Persons Ever				
of Diagnosis	Prevalence ¹	Prevalence	Living with	Living with	Diagnosed				
•		Rate ²	AIDS ³	HIV not AIDS ³	with AIDS				
Total Michigan	15,500	156.0	5,135	5,581	12,760				
Mackinac	10		1	0	2				
Macomb	570	72.3	184	186	477				
Manistee	20	81.5	6	4	9				
Marquette	20	30.9	4	9	17				
Mason	20	70.7	8	4	11				
Mecosta	20	49.3	4	7	11				
Menominee	10		0	4	3				
Midland	30	36.2	14	8	25				
Missaukee	10		1	5	3				
Monroe	60	41.1	28	13	64				
Montcalm	30	49.0	10	11	20				
Montmorency	10		2	0	3				
Muskegon	120	70.5	35	45	81				
Newaygo	20	41.8	8	5	22				
Oakland	1,650	138.2	507	558	1,162				
Oceana	10		3	4	9				
Ogemaw	10		1	1	3				
Ontonagon	10		0	1	0				
Osceola	10		3	2	8				
Oscoda	10		0	1	2				
Otsego	10		5	3	8				
Ottawa	100	42.0	38	27	93				
Presque Isle	10		0	0	2				
Roscommon	10	39.3	6	3	13				
Saginaw	200	95.2	63	68	156				
Sanilac	20		6	4	12				
Schoolcraft	10		1	1	3				
Shiawassee	30	65.9	11	7	18				
St. Clair	100	60.9	25	38	69				
St. Joseph	40	55.8	13	10	34				
Tuscola	20	34.3	6	5	14				
Van Buren	80	104.9	24	25	46				
Washtenaw	500	154.8	165	160	359				
Wayne	1,520	73.7	522	458	1,219				
City of Detroit	6,810	715.9	2,157	2,247	5,801				
Wexford	10		2	6	4				
PRISONS ⁴	700	N/A	220	472	482				
UNKNOWN	10	N/A	1	4	3				

^{1.} This estimate includes all persons living with HIV or AIDS, including those not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10.

^{2.} Rates are calculated per 100,000 population in 2000. Rates are unreliable for counties with the minimum estimated prevalence of 10, and are therefore not listed.

^{3.} Includes reports of HIV infection or AIDS that contain patient name or are otherwise unduplicated.

^{4.} The category PRISONS includes those persons who were in prison at the time of their HIV or AIDS diagnosis.

Table 3: Michigan Residents Reported Living with HIV or AIDS: Sex by Race by Behavior

							Othe	r or		
MALES:	White		Black		Hispanic		Unknown		TOTAL	
Male-Male Sex	2,517	74%	2,127	48%	134	45%	66	41%	4,844	59%
Injecting Drug Use	172	5%	773	17%	57	19%	6	4%	1,008	12%
Male-Male Sex/IDU	215	6%	326	7%	14	5%	6	4%	561	7%
Blood Recipient	81	2%	23	1%	1	0%	2	1%	107	1%
Heterosexual	85	3%	321	7%	30	10%	6	4%	442	5%
Perinatal	12	0%	53	1%	2	1%	0	0%	67	1%
Undetermined/Other	312	9%	801	18%	57	19%	76	47%	1,246	15%
Presumed Heterosexual	186	5%	558	13%	44	15%	24	15%	812	10%
Other	126	4%	243	5%	13	4%	52	32%	434	5%
MALE TOTAL	3,394	(41%)	4,424	(53%)	295	(4%)	162	(2%)	8,275	100%
							Other or			
FEMALES:	Whi	te	Black Hispanic		Unknown		TOTAL			
Injecting Drug Use	126	24%	522	29%	15	17%	6	13%	669	27%
Blood Recipient	11	2%	8	0%	0	0%	0	0%	19	1%
Heterosexual	269	51%	709	40%	52	59%	16	33%	1046	43%
Perinatal	11	2%	47	3%	5	6%	1	2%	64	3%
Undetermined/Other	112	21%	490	28%	16	18%	25	52%	643	26%
Presumed Heterosexual	97	18%	402	23%	15	17%	11	23%	525	22%
Other	15	3%	88	5%	1	1%	14	29%	118	5%
FEMALE TOTAL	529	(22%)	1,776	(73%)	88	(4%)	48	(2%)	2441	100%
GRAND TOTAL	3,923	37%	6,200	58%	383	4%	210	2%	10,716	100%

TECHNICAL NOTES

HIV Prevalence Estimate for Michigan

MDCH estimates that there are up to 15,500 HIV-infected persons (including those with AIDS) living in Michigan. This estimate includes all persons living in Michigan at diagnosis of HIV or AIDS, including those not reported or not yet diagnosed. It is supported by national estimates of HIV infection and rate of new AIDS diagnoses and deaths. Categorical estimates of HIV prevalence are calculated from the distribution of reported cases among each group of confidentially reported persons living with HIV or AIDS.

Footnotes for Table 1:

- 1. This estimate includes all persons living in Michigan at diagnosis of HIV or AIDS, including those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10.
- 2. Rates are calculated per 100,000 population in 2000.
- 3. Includes reports that contain patient name or are otherwise unduplicated.
- 4. Age, sex, race, and behavior percentages are calculated excluding missing data. The percentages of total cases missing this demographic information are given in parentheses.
- 5. The IDU risk category is further sub-divided to indicate the number and percentage of persons who also had a sexual partner who is considered to be a "high risk" heterosexual, (i.e., partner is an IDU, a bisexual male (for females), a recipient of HIV infected blood or blood products or a person who is known to be infected with HIV).
- 6. The heterosexual category includes only those persons with "high risk" heterosexual partners as defined in footnote 5.
- 7. This subset of undetermined includes persons who had heterosexual sex but their partner(s') risk is unknown. This includes one unconfirmed occupational exposure.
- 8. Includes persons with confirmed exposure in the health care setting in the U.S. (2) and pediatric cases with probable sexual mode of transmission (2).